

American plaice




Figure C1. Northeast Region American plaice has a controlled fishing mortality and biomass has not increasing as expected.

bigeye tuna - Atlantic




Figure C2. Highly Migratory Species bigeye tuna - Atlantic has a controlled fishing mortality but biomass is not increasing as expected. FMP not internationally implemented

ocean pout

Ratio

1

Overfishing $F/F_{msy} \geq 1$

Overfished $B/B_{msy} \leq 0.5$

0



1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005

Year

Overfished Declaration

Rebuilding Plan 2004-2014

Figure C3. Northeast Region ocean pout has a controlled fishing mortality but biomass has not increased as expected. B_{msy} proxy is in kg/tow.

sandbar shark




Figure C4. Highly Migratory Species sandbar shark has a controlled fishing mortality but biomass is not increasing as expected. FMP not internationally implemented

thorny skate




Figure C5. Northeast Region thorny skate has a controlled fishing mortality but biomass has not increased as expected. B_{msy} proxy is in kg/tow.

*Skate stocks F/F_{msy} are measured by percent change over a 3 year period.

windowpane flounder - SNE/MA




Figure C6. Northeast Region windowpane flounder – Southern New England / Mid-Atlantic Bight has a controlled fishing mortality but biomass has not increased as expected. B_{msy} proxy is in kg/tow.